



CALIFORNIA DEPARTMENT OF
FOOD & AGRICULTURE

Karen Ross, Secretary

FIFRA SECTION 2(ee) RECOMMENDATION FOR USE
IN THE STATE OF CALIFORNIA

CORETECT Tree and Shrub Tablets INSECTICIDE

EPA Reg. No. 432-1457

EPA SLN No. CA-080016

Use Directions for Glassy-winged sharpshooter (*Homalodisca vitripennis*)

It is a violation of federal law to use this product in any manner inconsistent with its labeling. This recommendation is permitted under FIFRA Section 2(ee). This Section 2(ee) label, the special local need label, and the federally registered label must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, and precautions on the EPA registered label.

DIRECTIONS FOR USE

For use in and around residential areas, industrial and commercial buildings.

CITRUS: Crops of group 10 including: calamondin, Citrus citron, citrus hybrids (includes chironja, tangelo, and tango), grapefruit, kumquat, lemon, lime, mandarin (tangerine), pummelo, orange (sweet and sour), Satsuma mandarin, white sapote (*Casimiroa* spp.), and other cultivars and hybrids of these.

ORNAMENTALS: *Murraya* spp. (Curry Tree, Orange Jessamine)

IN-GROUND

Pest Controlled	Coretect Tree and Shrub Tablets Insecticide
Glassy-winged sharpshooter (<i>Homalodisca vitripennis</i>)	2-3 tablets per inch of trunk diameter at breast height or per foot of shrub height

CONTAINERIZED

Pest Controlled	Coretect Tree and Shrub Tablets Insecticide
Glassy-winged sharpshooter (<i>Homalodisca vitripennis</i>)	1 tablet /ft ³ of container media

Restrictions:

Do not apply more than 103 tablets per 10,000 ft² per year. Preharvest interval (PHI): 0 days

Recommended applications:

Coretect Tree and Shrub Tablets Insecticide is formulated to provide consistent delivery of active ingredient over time. Release of active ingredient is dependent on the presence of adequate soil moisture. For best results, irrigate the plant after treatment, and ensure that the plant receives regular irrigation for at least 2-3 months after treatment.

Typically, the fine absorptive roots of trees are located near the base of the tree, in the upper six inches of the soil. Apply Coretect Tree and Shrub Insecticide Tablets near the base of the tree (i.e., within 6-24") to ensure rapid uptake by these absorptive roots. Space tablets evenly around the tree, 2-5" below the soil surface. Ensure that the tablets are completely covered with soil after the application.

For bushy trees and shrubs, numerous absorptive roots may be located near the drip-line. To apply Coretect Tree and Shrub Tablets Insecticide to plants with a definite drip-line, place tablets 2-5" below the soil surface, spaced evenly around the tree and along the drip-line.

